

Cardiovascular Risk in Hashimoto's Thyroiditis: Role of Thyroid Autoimmunity

Rajarajeswari Ranganadane¹, Sumathi Saravanakumar S², Srinivasan Abu Raghavan³, Asmathulla Shafiulla⁴, Girija Subramanian⁵, Maithili Karpaga Selvi Nachimuthu⁶

ABSTRACT

Hashimoto's thyroiditis (HT) is the most common autoimmune thyroid disease synonymous with hypothyroidism. The link between hypothyroidism and the risk of cardiovascular diseases is of contemporary interest. Studies have indicated the prevalence of metabolic syndrome and endothelial dysfunction in HT patients. HT *per se* might possess a role in atherosclerosis. Association of HT with dyslipidemia and chronic inflammation leading to endothelial dysfunction has been documented. However, the role of thyroid autoimmunity in promoting cardiovascular diseases remains unclear. Further studies unraveling the causal relationship between HT and cardiovascular disease would provide greater insight into the management of atherogenic complications observed in HT patients.

Keywords: Chronic inflammation, Dyslipidemia, Endothelial dysfunction, Hashimoto's thyroiditis.

SBV Journal of Basic, Clinical and Applied Health Science (2021): 10.5005/jp-journals-10082-03106

INTRODUCTION

Hashimoto's thyroiditis (HT) is an autoimmune disease characterized by the destruction of thyroid cells by both cell- and antibody-mediated immune responses¹ and is regarded as the commonest cause of hypothyroidism. HT has a prevalence rate of 1–4%. HT is more common in women and increases with age.^{2,3} Several studies have shown the association of hypothyroidism with dyslipidemia and chronic inflammation, which in turn enhances the risk for cardiovascular diseases.⁴ A marked increase in low-density lipoprotein (LDL) is seen in hypothyroidism, attributed to decreased LDL receptors in the liver culminating in reduced LDL clearance.^{5,6} Hyperlipidemia and chronic inflammation are implicated in atherosclerotic lesion formation. The levels of major inflammatory markers, such as interleukin (IL) 6, tumor necrosis factor-alpha (TNF- α), and high sensitive C-reactive protein (hs-CRP), were found to be elevated in HT patients.⁷ Though cardiovascular risk seen in overt hypothyroidism is a well-established fact, the influence of thyroid autoimmunity in increasing the cardiovascular events independent of thyroid dysfunction still remains a conundrum. Few studies suggest a putative role of thyroid autoimmunity in atherosclerosis development independent of thyroid dysfunction. Chronic inflammation and endothelial dysfunction could be the link. Cytokines are believed to modify epithelium thereby allowing infiltration of the thyroid by immune cells: a mechanism underlying HT.⁸ Thus, the purpose of this review is to primarily provide the discerning researchers an overall picture of cardiovascular risk seen in HT patients, which might be a great source of help in the management.

DYSLIPIDEMIA AND HASHIMOTO'S THYROIDITIS: THE NEXUS

Dyslipidemia as observed in thyroid abnormalities is a potent risk factor of cardiovascular events among patients with abnormal thyroid function.⁹ Studies have shown a positive association of extracellular thyroid stimulating hormone (TSH) with lipid levels in hypothyroid subjects. In a prominent study referred to as the HUNT study, the association of hypothyroidism with high blood lipids was

^{1,4}Department of Biochemistry, Sri Manakula Vinayagar Medical College and Hospital, Puducherry, India

^{2,3}Department of Biochemistry, Mahatma Gandhi Medical College and Research Institute, Puducherry, India

⁵Department of General Medicine, Sri Manakula Vinayagar Medical College and Hospital, Puducherry, India

⁶Department of Biochemistry, All India Institute of Medical Sciences, Jodhpur, India

Corresponding Author: Sumathi Saravanakumar, Department of Biochemistry, Mahatma Gandhi Medical College and Research Institute, Puducherry, India, e-mail: sumathi.bio21@gmail.com

How to cite this article: Ranganadane R, Sumathi S, Raghavan SA, Shafiulla A, Subramanian G, Nachimuthu MKS. Cardiovascular Risk in Hashimoto's Thyroiditis: Role of Thyroid Autoimmunity. *J Basic Clin Appl Health Sci* 2021;4(1):20–22.

Source of support: Nil

Conflict of interest: None

observed as a linear function across the entire reference range of TSH.¹⁰ Tagami et al. determined an association between serum TSH and lipid levels.¹¹ A pronounced atherosclerotic lesion has been demonstrated in children and adolescents with hypothyroidism, with an accompanying elevation in total cholesterol (TC) and LDL levels and decreased high-density lipoprotein (HDL) level.¹²

Endothelial Dysfunction—An Important Point for Consideration

Whickham's study showed no association between elevated TSH and dyslipidemia.¹³ A study by Tamer et al. also showed no discernible correlation of TSH levels with serum lipids. However, an interesting point had emerged, viz., anti-thyroperoxidase antibodies (TPOAb) have correlated positively and pronouncedly with serum triglyceride (TG).¹⁴ Increased carotid intima-media thickness in HT women, independent of thyroid function has also been reported.¹⁵ Subclinical hypothyroid patients are more prone to

developing atherosclerotic cardiovascular disease, in spite of having lower TC levels when compared to healthy controls.¹⁶ As per another study, no pronounced difference in serum lipid levels between the hypothyroid and healthy controls was observed.^{17,18} According to Volpe's hypothesis,¹⁹ due to defective suppressor T cells, T cells are not suppressed and produce a lot of cytokines such as IFN- γ , IL-2, and TNF- α . These cytokines might cause hyperlipidemia as suggested by another study.²⁰ Hyperlipidemia diminishes the expression of endothelial nitric oxide synthase (eNOS) and increases asymmetric dimethylarginine levels, an eNOS endogenous inhibitor, ultimately leading to endothelial dysfunction.^{21,22}

Inflammatory Markers and Hashimoto's Thyroiditis

Inflammation is another probable factor that could cause endothelial dysfunction. Acute and chronic inflammations are strongly related to endothelial dysfunction.²³⁻²⁵ A study done by Türemen et al. has shown an increase in inflammatory factors, such as IL-6, hs-CRP, and TNF- α , in HT patients. Furthermore, these are positively correlated with flow-mediated dilation. This event points to the contribution of low-grade chronic inflammation in promoting atherosclerosis in HT subjects.⁷ IL-6 plays a pivotal role in vascular inflammation by promoting endothelial dysfunction, smooth muscle cell proliferation, and migration.²⁵ Few studies documenting an increase in serum IL-6 levels in HT are available.^{26,27}

hs-CRP, a marker of systemic inflammation, is used as a biomarker for cardiovascular risk assessment. A point of significance is that endothelial dysfunction attributed to dyslipidemia and low-grade chronic inflammation is purported to be the contributing factors for cardiovascular disorders visualized in HT.⁷ However, the role of thyroid autoimmunity is unclear. According to another study, no correlation between anti-TPO and IL-6 or anti-TPO and IL-15 in HT subjects irrespective of thyroid function status was observed.²⁸ The exact role of antibodies against thyroid peroxidase remains a mystery and it is suggested that they may promote the release of cytokines, including IL-6, TNF- α , and INF- γ .²⁹ Few studies describe a correlation between TSH and TPOAb that might promote the release of cytokines, including IL-6, TNF- α , and INF- γ .^{29,30} TSH may promote endothelial dysfunction by increasing IL-6 and TNF- α secretion by binding to adipocyte TSH receptor and bone marrow cell TSH receptor, respectively.⁴ Higher levels of

TSH could be proinflammatory stimuli of adipocytes.³¹ In a study done by Sieminska et al., thyroid autoimmunity was found to be associated with elevated IL-6 concentration, regardless of thyroid function. However, no association was found between TSH and IL-6 levels.⁸ However, in the analysis done by LeGrys et al., there was no connection between the presence of thyroid antibodies and the risk of myocardial infarction (Flowchart 1).³²

Taddei et al. observed an increase in IL-6 and hs-CRP levels in HT subjects, which could trigger endothelial dysfunction, one of the early mechanisms promoting atherosclerosis and cardiovascular disease. IL-6 promotes atherosclerosis either directly by endothelial-dependent mechanisms or indirectly by stimulating hepatic production of hs-CRP.²⁷ CRP may interfere with endothelial function by downregulating eNOS and by upregulation of endothelin-1, a potent vasoconstrictor that antagonizes nitric oxide action.³³ Though the clear association of elevated CRP with HT subjects is shown in certain studies,²⁷ Pearce et al. could not find any difference in CRP levels among HT patients and controls.³⁴


CONCLUSION

Thyroid autoimmunity is found to be associated with dyslipidemia and low-grade chronic inflammation, which might result in future cardiovascular morbidities. However, there is significant heterogeneity of results among the studies included. Further studies are needed to address the causality of cardiovascular diseases in HT patients, which may have implications in understanding the disease mechanisms and future treatment strategies.

REFERENCES

1. Yang M, Du C, Wang Y, Liu J. CD19+CD24hiCD38hi regulatory B cells are associated with insulin resistance in type I Hashimoto's thyroiditis in Chinese females. *Exp Ther Med* 2017;14(4):3887–3893. DOI: 10.3892/etm.2017.4925.
2. Cunha CA, Neves C, Neves JS, Oliveira SC, Sokhatska O, Dias C, et al. Cardiovascular risk factors in patients with autoimmune thyroiditis. *Rev Port Endocrinol Diabetes Metab* 2017;12(2):133–141. DOI:10.1530/endoabs.49.gp199.
3. Sood N, Nigam JS. Correlation of fine needle aspiration cytology findings with thyroid function test in cases of lymphocytic thyroiditis. *J Thyroid Res* 2014;2014:430510. DOI:10.1155/2014/430510.
4. Lu M, Yang CB, Gao L, Zhao JJ. Mechanism of subclinical hypothyroidism accelerating endothelial dysfunction (review). *Exp Ther Med* 2015;9(1):3–10. DOI: 10.3892/etm.2014.2037.
5. Duntas LH. Thyroid disease and lipids. *Thyroid* 2002;12(4):287–293. DOI: 10.1089/10507250252949405.
6. Farhangi MA, Eshraghian M, Keshavarz SA, Saboor-Yaraghi AA. Thyroid stimulating hormone, triiodothyronine and thyroxine concentrations and their relationship with metabolic parameters, anthropometric variables and body composition in premenopausal euthyroid obese women. *Turk J Endocrinol Metab* 2015;19(1):1–6. DOI: 10.4274/tjem.2562.
7. Türemen EE, Çetinarslan B, Şahin T, Cantürk Z, Tarkun İ. Endothelial dysfunction and low grade chronic inflammation in subclinical hypothyroidism due to autoimmune thyroiditis. *Endocr J* 2011;58(5):349–354. DOI: 10.1507/endocrj.k10e-333.
8. Siemińska L, Wojciechowska C, Walczak K, Borowski A, Marek B, Nowak M, et al. Associations between metabolic syndrome, serum thyrotropin, and thyroid antibodies status in postmenopausal women, and the role of interleukin-6. *Endokrynol Pol* 2015;66(5):394–403. DOI: 10.5603/EP.2015.0049.
9. Farhangi MA, Dehghan P, Tajmiri S. Powdered black cumin seeds strongly improves serum lipids, atherogenic index of plasma and modulates anthropometric features in patients with Hashimoto's thyroiditis. *Lipids Health Dis* 2018;17(1):59. DOI: 10.1186/s12944-018-0704-x.

Flowchart 1: Role of thyroid autoimmunity in cardiovascular disease manifestation in Hashimoto's thyroiditis



10. Asvold BO, Vatten LJ, Nilsen TI, Bjørø T. The association between TSH within the reference range and serum lipid concentrations in a population-based study. The HUNT study. *Eur J Endocrinol* 2007;156(2):181–186. DOI: 10.1530/eje.1.02333.
11. Tagami T, Tamanaha T, Shimazu S, Honda K, Nanba K, Nomura H, et al. Lipid profiles in untreated patients with Hashimoto's thyroiditis and the effects of thyroxine treatment on subclinical hypothyroidism with Hashimoto's thyroiditis. *Endocr J* 2010;57(3):253–258. DOI: 10.1507/endocrj.k09e-315.
12. Rodrigues AN, Abreu GR, Resende RS, Goncalves WL, Gouvea SA. Cardiovascular risk factor investigation: a pediatric issue. *Int J General Med* 2013;6:57–66. DOI: 10.2147/IJGM.S41480.
13. Tunbridge WM, Evered DC, Hall R, Appleton D, Brewis M, Clark F, et al. Lipid profiles and cardiovascular disease in the Whickham area with particular reference to thyroid failure. *Clin Endocrinol (Oxf)* 1977;7(6):495–508. DOI: 10.1111/j.1365-2265.1977.tb01341.x.
14. Tamer G, Mert M, Tamer I, Mesci B, Kiliç D, Arik S. Effects of thyroid autoimmunity on abdominal obesity and hyperlipidemia. *Pol J Endocrinol* 2011;62(5):421–428. PMID: 22069103
15. Topaloglu O, Gokay F, Kucukler K, Burnik FS, Mete T, Yavuz HC, et al. Is autoimmune thyroiditis a risk factor for early atherosclerosis in premenopausal women even if in euthyroid status? *Endocrine* 2013;44(1):145–151. DOI: 10.1007/s12020-012-9842-5.
16. Hak AE, Pols HA, Visser TJ, Drexhage HA, Hofman A, Witteman JC. Subclinical hypothyroidism is an independent risk factor for atherosclerosis and myocardial infarction in elderly women: the Rotterdam Study. *Ann Intern Med* 2000;132(4):270–278. DOI: 10.7326/0003-4819-132-4-200002150-00004.
17. Uzunlulu M, Yorulmaz E, Oguz A. Prevalence of subclinical hypothyroidism in patients with metabolic syndrome. *Endocrine J* 2007;54(1):71–76. DOI: 10.1507/endocrj.k06-124.
18. Shantha GP, Kumar AA, Jeyachandran V, Rajamanickam D, Rajkumar K, Salim S, et al. Association between primary hypothyroidism and metabolic syndrome and the role of C reactive protein: a cross-sectional study from South India. *Thyroid Res* 2009;2(1):2. DOI: 10.1186/1756-6614-2-2.
19. Volpe R. Autoimmune thyroiditis. In: Braverman LE, Utiger RD, editors. *Werner and Ingbar's The Thyroid*. Philadelphia: JB Lippincott Co.;1991. p. 921–941.
20. Sultan A, Strodtloff D, Robertson AK, Paulsson-Berne G, Fauconnier J, Parini P, et al. T cell-mediated inflammation in adipose tissue does not cause insulin resistance in hyperlipidemic mice. *Circ Res* 2009;104(8):961–968. DOI: 10.1161/CIRCRESAHA.108.190280.
21. Laufs U, La Fata V, Plutzky J, Liao JK. Upregulation of endothelial nitric oxide synthase by HMG CoA reductase inhibitors. *Circulation* 1998;97(12):1129–1135. DOI: 10.1161/01.cir.97.12.1129.
22. Ito A, Tsao PS, Adimoolam S, Kimoto M, Ogawa T, Cooke JP. Novel mechanism for endothelial dysfunction: dysregulation of dimethylarginine dimethylaminohydrolase. *Circulation* 1999;99(24):3092–3095. DOI: 10.1161/01.cir.99.24.3092.
23. Hingorani AD, Cross J, Kharbanda RK, Mullen MJ, Bhagat K, Taylor M, et al. Acute systemic inflammation impairs endothelium-dependent dilatation in humans. *Circulation* 2000;102(9):994–999. DOI: 10.1161/01.cir.102.9.994.
24. Fichtlscherer S, Rosenberger G, Walter DH, Breuer S, Dimmeler S, Zeiher AM. Elevated C-reactive protein levels and impaired endothelial vasoreactivity in patients with coronary artery disease. *Circulation* 2000;102(9):1000–1006. DOI: 10.1161/01.cir.102.9.1000.
25. Kharbanda RK, Walton B, Allen M, Klein N, Hingorani AD, MacAllister RJ, et al. Prevention of inflammation-induced endothelial dysfunction: a novel vasculo-protective action of aspirin. *Circulation* 2002;105(22):2600–2604. DOI: 10.1161/01.cir.0000017863.52347.6c.
26. Bai X, Sun J, Wang W, Shan Z, Zheng H, Li Y, et al. Increased differentiation of Th22 cells in Hashimoto's thyroiditis. *Endocr J* 2014;61(12):1181–1190. DOI: 10.1507/endocrj.EJ14-0265.
27. Taddei S, Caraccio N, Virdis A, Dardano A, Versari D, Ghiadoni L, et al. Low-grade systemic inflammation causes endothelial dysfunction in patients with Hashimoto's thyroiditis. *J Clin Endocrinol Metab* 2006;91(12):5076–5082. DOI: 10.1210/jc.2006-1075.
28. Elshenawy SZ, Hemi MH, Attia H. Serum levels of pro-inflammatory cytokines (interleukin 6 and interleukin 15) and adiponectin in Hashimoto's thyroiditis with different thyroid function states. *J Am Sci* 2011;7(6):1156–1162. ISSN: 1545-1003
29. Nielsen CH, Brix TH, Leslie RGQ, Hegedüs L. A role for autoantibodies in enhancement of pro-inflammatory cytokine responses to a self-antigen, thyroid peroxidase. *Clin Immunol* 2009;133(2):218–227. DOI: 10.1016/j.clim.2009.07.014.
30. Sieminska L, Wojciechowska C, Kos-Kudla B, Marek B, Kajdaniuk D, Nowak M, et al. Serum concentrations of leptin, adiponectin, and interleukin-6 in postmenopausal women with Hashimoto's thyroiditis. *Endokrynol Pol* 2010;61(1):112–116. PMID: 20205113
31. Gagnon A, Langille ML, Chaker S, Antunes TT, Durand J, Sorisky A. TSH signaling pathways that regulate MCP-1 in human differentiated adipocytes. *Metabolism* 2014;63(6):812–821. DOI: 10.1016/j.metabol.2014.02.015.
32. LeGrys VA, Funk MJ, Lorenz CE, Giri A, Jackson RD, Manson JE, et al. Subclinical hypothyroidism and risk for incident myocardial infarction among postmenopausal women. *J Clin Endocrinol Metab* 2013;98(6):2308–2317. DOI: 10.1210/jc.2012-4065.
33. Yu YT, Ho CT, Hsu HS, Li CI, Davidson LE, Liu CS, et al. Subclinical hypothyroidism is associated with elevated high-sensitive C-reactive protein among adult Taiwanese. *Endocrine* 2013;44(3):716–722. DOI: 10.1007/s12020-013-9915-0.
34. Pearce EN, Bogazzi F, Martino E, Brogioni S, Pardini E, Pellegrini G, et al. The prevalence of elevated serum C-reactive protein levels in inflammatory and non-inflammatory thyroid disease. *Thyroid* 2003;13(7):643–648. DOI: 10.1089/105072503322239989.